

17-1074  
12-1074  
000  
93113

**Multiple Spectral Components in Nearby Radio Galaxies**

**NASA Grant NAG5-2791**

**Annual Reports No's 1 and 2**

**For the Period 1 October 1994 through 30 September 1996**

**Principal Investigator  
Dr. Mark Birkinshaw**

**September 1996**

**Prepared for:**

**National Aeronautics and Space Administration  
Goddard Space Flight Center  
Greenbelt, Maryland 20771**

**Smithsonian Institution  
Astrophysical Observatory  
Cambridge, Massachusetts 02138**

**The Smithsonian Astrophysical Observatory  
is a member of the  
Harvard-Smithsonian Center for Astrophysics**

The NASA Technical Officer for this grant is Dr. Nicholas White, Code 668, Laboratory for High Energy Astrophysics, NASA/Goddard Space Flight Center, Greenbelt, MD 20771.

The ASCA observation of NGC 6251 returned significant spectral data, and are showing that the interpretation of NGC 6251 which we based on earlier ROSAT spectra needs modification.

A simple power-law fit to NGC 6251, using all the GIS and SIS data, produces a good reduced chi-squared (0.97; 749 degrees of freedom), but between 1 and 2 keV there is a significant pattern of residuals in both detectors which suggests that the fit is imperfectly describing the spectrum.

A superposition of a power law spectrum and a thermal spectrum, which was the model most favored by our ROSAT data, produces an adequate (but worse) fit, with reduced chi-squared 1.03 (745 degrees of freedom), and only small changes in chi-squared are obtained as the abundance varies.

An excellent fit (reduced chi-squared = 0.94, 747 degrees of freedom) is produced by an alternative model based on the ROSAT data, where two thermal components make up the X-ray spectrum. The pattern of residuals is much superior to that found with the power-law spectrum. If the abundances are allowed to fall to 0.3 cosmic, then even better fits are produced (reduced chi-squared of 0.90)

All these fits require absorption above the galactic value, and the best of them appear to involve both a cool and a hot component of thermal gas (though the hot component can be replaced by power-law emission with only a modest cost in goodness of fit). The excess absorption is consistent with the neutral hydrogen column that we found with the VLA, so that a consistent picture of a cool atmosphere and an absorbing column near the central source is emerging. What is not clear at the moment is whether the central object is principally non-thermal emission (a mini-AGN) or thermal emission (a hot bubble or corona of a disc or cloud).

Further comparisons of the X-ray data from ASCA and ROSAT, including joint spectral fitting and absorption data from the VLA studies, may resolve this scientific issue. We also plan checks of the result using a wider variety of fitting codes, since there are well-known problems in interpreting structures in X-ray spectra due to model uncertainties in the line emission from atomic species in astrophysical objects.